

Products of characters and derived length II

by

Edith Adan-Bante

Department of Mathematics

University of Illinois at Urbana-Champaign

Urbana, IL 61801 USA

Currently at the University of Southern Mississippi Gulf Coast, 730 East
Beach Blvd, Long Beach, MS 39560

The author was partially supported by the National Science Foundation
by grant DMS 99-70030.

1 Introduction

Let G be a finite group. Let χ and ψ be irreducible complex characters, i.e. irreducible characters over the complex numbers. Since a product of characters is a character, $\chi\psi$ is a character. Then the decomposition of the character $\chi\psi$ into its distinct irreducible constituents $\alpha_1, \alpha_2, \dots, \alpha_n$ has the form

$$\chi\psi = \sum_{i=1}^n a_i \alpha_i$$

where $n > 0$ and $a_i > 0$ is the multiplicity of α_i for each $i = 1, \dots, n$. Let $\eta(\chi\psi) = n$ be the number of distinct irreducible constituents of the character $\chi\psi$. Denote by $\text{Ker}(\chi\psi)$ the kernel of the character $\chi\psi$. Also denote by $\text{dl}(\text{Ker}(\alpha)/\text{Ker}(\chi\psi))$ the derived length of the group $\text{Ker}(\alpha)/\text{Ker}(\chi\psi)$.

Given $\chi \in \text{Irr}(G)$, let $\bar{\chi}$ be the complex conjugate of χ , i.e. $\bar{\chi}(g) = \overline{\chi(g)}$ for all $g \in G$. In Theorem A of [1], it is proved that there exist universal constants C_0 and D_0 such that for any finite solvable group G and any $\chi \in \text{Irr}(G)$ we have that $\text{dl}(G/\text{Ker}(\chi)) \leq C_0 \eta(\chi\bar{\chi}) + D_0$. The following generalizes that result and it is the main result of this paper.

Theorem A. *There exist universal constants C and D such that for any finite solvable group G , any irreducible characters χ and ψ of G , and any irreducible constituent α of $\chi\psi$, we have*

$$\text{dl}(\text{Ker}(\alpha)/\text{Ker}(\chi\psi)) \leq C \eta(\chi\psi) + D.$$

In Theorem 2.8 we prove that we may take $C = 2$ and $D = -1$ if the group G in Theorem A is, in addition, supersolvable.

Theorem B. *Let G be a finite group and $\chi, \psi \in \text{Irr}(G)$ be characters. If $(\chi(1), \psi(1)) = 1$, then $\text{dl}(\text{Ker}(\alpha)/\text{Ker}(\chi\psi)) \leq 1$ for any irreducible constituent α of the product $\chi\psi$.*

Thus if $(\chi(1), \psi(1)) = 1$, then the irreducible constituents of the product $\chi\psi$ are “almost” faithful characters of the group $G/\text{Ker}(\chi\psi)$.

Acknowledgment. I thank Professor Everett C. Dade for helpful discussions. I also thank the referee for her/his suggestions for improving the presentation of this work, and especially for shortening the proof of Theorem A.

2 Proof of Theorem A

We will be using the notation of [2]. In addition, we write $M \subseteq N$ if M is a subgroup of N . Also $M \subset N$ denotes that M is a proper subgroup of N .

Definition 2.1. Assume that G acts on a finite vector space V . We define $m(G, V)$ as the number of orbits of nonzero vectors under the action of G .

The key tool for the proof of Theorem A is the following.

Lemma 2.2 (Keller). *There exist universal constants A and B such that for any finite solvable group G acting faithfully on a finite vector space V we have that*

$$\text{dl}(G) \leq Am(G, V) + B.$$

Proof. See [3]. □

We emphasize that the result of Keller is much stronger than Lemma 2.2. He proved that the derived length of G is bounded by a logarithmic function of the number of different sizes of G -orbits on V .

Definition 2.3. We define the function f by

$$f(n) = An + B,$$

for any integer $n \geq 1$, where A and B are as in Lemma 2.2.

Definition 2.4. Let G be a finite group and M and N be normal subgroups of G such that $M \subset N$. Let $\theta \in \text{Irr}(N)$. We say that (N, M, θ) is an **extreme triple** of G if for any normal subgroup K of G such that $M \subset K \subseteq N$, we have that θ_K is irreducible but θ_M is reducible.

Observe that given $N \trianglelefteq G$ and a nonlinear character $\theta \in \text{Irr}(N)$ of N , we can always find $M \triangleleft G$ such that (N, M, θ) is an extreme triple.

Lemma 2.5. *Let (N, M, θ) be an extreme triple of G and suppose $M \subset L \subseteq N$, where $L \triangleleft G$ and L/M is abelian. Then $\mathbf{C}_{N/M}(L/M)$ is abelian.*

Proof. By definition of extreme triple, we have that $\theta_L \in \text{Irr}(L)$ and θ_M is reducible. Since L/M is abelian, by Theorem 6.22 of [2] we have that there exists a subgroup U containing M such that U has prime index in L and θ_U is reducible. Since θ_U is reducible and $|L : U| = p$, we have that θ_U is a sum of p distinct irreducible characters of U . Let $\varphi \in \text{Irr}(U)$ be an irreducible constituent of θ_U .

Let $C = \mathbf{C}_N(L/M)$. Note that $L \subseteq C$ and thus $U \triangleleft C$ and $\theta_C \in \text{Irr}(C)$. It follows that $|C : C_\varphi| = p$ and $C = LC_\varphi$. Since L centralizes C/M , we see that $C_\varphi \trianglelefteq C$ and C/C_φ is abelian. Then $MC' \subseteq C_\varphi$ and thus $\theta_{MC'}$ is reducible. Since $MC' \trianglelefteq G$, it follows that $MC' = M$. Thus $C' \subseteq M$ as wanted. $\square$

We introduce more notation. If N/M is a normal section of G and Δ is a character of G , we write $S_\Delta(N/M)$ to denote the (possibly empty) set of those irreducible constituents α of Δ such that $M \subseteq \text{Ker}(\alpha)$ and $N \not\subseteq \text{Ker}(\alpha)$.

Lemma 2.6. *Let (N, M, θ) be an extreme triple of G , where N/M is solvable. Assume that $\theta\bar{\theta}$ is (not necessarily irreducible) constituent of Δ_N , for some character Δ of G . Then $S_\Delta(N/M)$ is nonempty, and the derived length of N/M is at most $1 + f(|S_\Delta(N/M)|)$, where f is the function as in Definition 2.3.*

Proof. Since N/M is solvable and non-trivial, we can choose $L \trianglelefteq G$ with $M \subset L \subseteq N$ such that L/M is a chief factor of G . Also, as in the previous lemma, we know that there exists a subgroup U containing M and of prime index p in L such that θ_U is a sum of p distinct irreducible constituents. Thus θ vanishes on $L \setminus U$. Since θ_L is N -invariant, θ vanishes on $L \setminus D$, where D is the intersection of the N -conjugates of U . Since $M \subseteq D \subset L$ and $D \triangleleft N$, we can choose a chief factor L/K of N such that $D \subseteq K$, and thus θ vanishes on $L \setminus K$.

Write $\varphi = \theta_L \in \text{Irr}(L)$ and let λ be a nonprincipal linear character of L/K . Then $\varphi\lambda = \varphi$, and this λ is a constituent of $\varphi\bar{\varphi}$, which in turn is a constituent of Δ_L . It follows that λ is a constituent of α_L , where α is an irreducible constituent of Δ . In particular, we see that $\alpha \in S_\Delta(N/M)$, which is therefore nonempty.

Set $H = \mathbf{N}_G(K)$. Observe that $H \supseteq N$. All members of the H -orbit of λ are linear constituents of α_L having kernels containing K . Conversely, we show that if ν is any linear constituent of α_L such that $K \subseteq \text{Ker}(\nu)$, then ν lies in the H -orbit of λ . Certainly $\nu = \lambda^g$ for some element $g \in G$, and thus ν is nonprincipal. Also, since $K \subseteq \text{Ker}(\lambda)$, we see that $K^g \subseteq \text{Ker}(\nu)$. We assumed that $K \subseteq \text{Ker}(\nu)$, and thus $KK^g \subseteq \text{Ker}(\nu) \subset L$. Observe that $KK^g \triangleleft N$ since $K^g \subseteq N^g = N$ and K is normal in N . Since $KK^g \triangleleft N$ and L/K is a chief factor of N , it follows that $K = K^g$ and thus $g \in H$. We see

now that for each H -orbit of nonprincipal linear characters of L/K , there is a member of $S_\Delta(N/M)$ that lies over all members of this orbit and over no other linear character of L/K . It follows that the number of H -orbits on L/K , which is the same as the number of H -orbits of linear characters of L/K , is at most $|S_\Delta(N/M)|$. By Lemma 2.2, therefore, the derived length of $H/\mathbf{C}_H(L/K)$ is at most $f(|S_\Delta(N/M)|)$. Set $m = f(|S_\Delta(N/M)|)$. Then the derived length of $N/\mathbf{C}_N(L/K)$ is at most m and thus $N^{(m)} \subseteq \mathbf{C}_N(L/K)$. Since $[L, N^{(m)}] \subseteq K$, $K \subset L$, L/M is a chief factor of G and $[L, N^{(m)}] \triangleleft G$, we see that $[L, N^{(m)}] \subseteq M$. Thus $N^{(m)}$ centralizes L/M . Since $\mathbf{C}_{N/M}(L/M)$ is abelian by the previous lemma, it follows that $N^{(m+1)} \subseteq M$, as desired. $\square$

Lemma 2.7. *Let G be a finite group and $\chi, \psi \in \text{Irr}(G)$. Let $\alpha \in \text{Irr}(G)$ be an irreducible constituent of the product $\chi\psi$ and $K = \text{Ker}(\alpha)$. Then there exists some $\theta \in \text{Irr}(K)$ such that $[\chi_K, \theta] \neq 0$ and $[\overline{\psi}_K, \theta] \neq 0$.*

Proof. Since $K = \text{Ker}(\alpha)$, we have that $\alpha_K = \alpha(1)1_K$. Thus

$$\alpha(1)[(\chi\psi)_K, 1_K] = [(\chi\psi)_K, (\alpha)_K] > 0.$$

Therefore there exist $\theta, \sigma \in \text{Irr}(K)$ such that $[\chi_K, \theta] \neq 0$, $[\psi_K, \sigma] \neq 0$ and $[\theta\sigma, 1_K] \neq 0$. Since θ and σ are irreducible characters and $[\theta\sigma, 1_K] = [\theta, \overline{\sigma}] > 0$, we conclude that $\theta = \overline{\sigma}$. Thus $[\overline{\psi}_K, \theta] \neq 0$. $\square$

Proof of Theorem A. Write $\Delta = \chi\psi$ and $n = \eta(\chi\psi)$. Note that we can assume that Δ is faithful. Let $K = \text{Ker}(\alpha)$ so that our task is to show that the derived length $\text{dl}(K)$ is bounded by a linear function on n . By Lemma 2.7, we can choose a character $\nu \in \text{Irr}(K)$ that lies under both χ and $\overline{\psi}$. Also we see that if $N \triangleleft G$ and $N \subseteq \text{Ker}(\nu)$, then $N \subseteq \text{Ker}(\Delta)$ and thus $N = 1$.

Write $K_0 = K$. If ν is nonlinear, we can choose a subgroup L such that (K, L, ν) is an extreme triple of G . Write $K_1 = L$ and let ν_1 be an irreducible constituent of ν_L . If ν_1 is nonlinear, we can repeat the process and choose an extreme triple of G of the form (K_1, K_2, ν_1) . Continuing like this, we obtain a series $K = K_0 \supset K_1 \supset \cdots \supset K_r$ of normal subgroups of G and characters $\nu_i \in \text{Irr}(K_i)$ such that (K_i, K_{i+1}, ν_i) is an extreme triple for $0 \leq i < r$ and where ν_r is linear. Also, we know that for each subscript i , the character ν_i is a constituent of ν_{K_i} , and thus ν_i lies under both χ and $\overline{\psi}$. It follows that $\nu_i \overline{\nu_i}$ is a constituent of Δ_{K_i} . Therefore by Lemma 2.6, the sets $S_\Delta(K_i/K_{i+1})$ are nonempty for $0 \leq i < r$. Since they are certainly disjoint and α lies in none of the sets $S_\Delta(K_i/K_{i+1})$, we have that $r < n$. Also, writing $s_i = |S_\Delta(K_i/K_{i+1})|$, we see that $\sum s_i < n$.

By Lemma 2.6, the derived length of K_i/K_{i+1} is at most $1 + f(s_i)$. Also, because ν_r is linear, we see that $(K_r)' \subseteq \text{Ker}(\nu_r)$. Since $(K_r)' \triangleleft G$, it follows that $(K_r)' \subseteq \text{Ker}(\Delta)$, and hence $(K_r)' = 1$ and K_r is abelian. We see now that the derived length $\text{dl}(K)$ is at most $1 + \sum_{i=0}^{r-1} (1 + f(s_i))$. Since $f(x) = Ax + B$, we have that $\text{dl}(K) \leq A \sum s_i + rB + r + 1$. But $\sum s_i < n$ and $r < n$, and so $\text{dl}(K) < (A + B + 1)n + 1$. The proof is now complete. $\square$

Theorem 2.8. *Let G be a finite supersolvable group and $\chi\psi \in \text{Irr}(G)$ be characters of G . Let $\alpha \in \text{Irr}(G)$ be any constituent of the product $\chi\psi$. Then*

$$\text{dl}(\text{Ker}(\alpha)/\text{Ker}(\chi\psi)) \leq 2\eta(\chi\psi) - 1.$$

Proof. Set $K = \text{Ker}(\alpha)$ and $n = \eta(\chi\psi)$. We choose a character $\nu \in \text{Irr}(K)$ that lies under both χ and $\overline{\psi}$. As in the proof of Theorem A, we obtain a series $K = K_0 \supset K_1 \supset \cdots \supset K_r$ of normal subgroups of G and characters $\nu_i \in \text{Irr}(K_i)$ such that (K_i, K_{i+1}, ν_i) is an extreme triple for $0 \leq i < r$

and where ν_r is linear. As before, we have that $r < n$ and $K_r/\text{Ker}(\chi\psi)$ is abelian.

For $0 \leq i < r$, let $L_i \triangleleft G$ such that $K_i \subset L_i \subseteq K_{i_1}$ and L_i/K_i is a chief factor of G . Since G is supersolvable, we have that L_i/K_i is cyclic of prime order and therefore $G/\mathbf{C}_G(L_i/K_i)$ is abelian. Thus $\text{dl}(K_i/K_{i+1}) \leq 2$ for $0 \leq i < r$. Therefore

$$\text{dl}(K/\text{Ker}(\chi\psi)) \leq \sum_{i=1}^r \text{dl}(K_{i-1}/K_i) + \text{dl}(K_r/\text{Ker}(\chi\psi)) \leq 2r + 1.$$

Since $r < n$, we conclude that $\text{dl}(K/\text{Ker}(\chi\psi)) \leq 2(n-1) + 1 = 2n-1$ and the proof is complete. $\square$

3 Proof of Theorem B

Proof of Theorem B. Set $K = \text{Ker}(\alpha)$. Let $\theta \in \text{Irr}(K)$ be as in Lemma 2.7. Since $[\chi_K, \theta] \neq 0$ and $[\overline{\psi_K}, \theta] \neq 0$, and K is normal in G , we have that $\theta(1)$ divides both $\chi(1)$ and $\psi(1)$. Since $(\chi(1), \psi(1)) = 1$, it follows that $\theta(1) = 1$.

Observe that $[K, K] \subseteq \text{Ker}(\theta) \subseteq K$ since $\theta(1) = 1$. Also observe that $[K, K]$ is normal in G since K is normal in G . Since θ is an irreducible constituent of χ , $[K, K] \subseteq \text{Ker}(\theta)$ and $[K, K]$ is normal in G , we have that $\chi_{[K, K]} = \chi(1)1_{[K, K]}$. Similarly we can check that $\psi_{[K, K]} = \psi(1)1_{[K, K]}$. Thus $[K, K] \subseteq \text{Ker}(\chi\psi) \subseteq K$. Therefore $K/\text{Ker}(\chi\psi)$ is abelian and thus $\text{dl}(\text{Ker}(\alpha)/\text{Ker}(\chi\psi)) \leq 1$. $\square$

Example 3.1. Let G be a solvable group and $\chi, \psi \in \text{Irr}(G)$. Assume that

$$\text{dl}(\text{Ker}(\alpha)/\text{Ker}(\chi\psi)) \leq 1$$

for all irreducible constituents α of the product $\chi\psi$. Then we do not necessarily have that $(\chi(1), \psi(1)) = 1$.

Proof. Let p be an odd prime. Let E be an extra-special group of order p^3 . Choose a character $\chi \in \text{Irr}(E)$ such that $\chi(1) = p$. Let $\psi = \overline{\chi}$. Observe that $\text{Ker}(\chi\psi) = \mathbf{Z}(E)$. Also observe that $(\chi(1), \psi(1)) = \chi(1) = p$. Since $E/\mathbf{Z}(E)$ is abelian, for all the irreducible constituents α of the product $\chi\psi$, we have that $\text{dl}(\text{Ker}(\alpha)/\text{Ker}(\chi\psi)) = 1$. $\square$

4 Further results

Assume that G is a finite group and χ, ψ are irreducible characters of G such that the product $\chi\psi$ has a linear constituent α . Observe that $\alpha\overline{\chi} \in \text{Irr}(G)$ since $\alpha(1) = 1$ and $\chi \in \text{Irr}(G)$. Since $[\psi, \alpha\overline{\chi}] = [\chi\psi, \alpha] > 0$, we have that $\psi = \alpha\overline{\chi}$. In particular $\chi(1) = \psi(1)$. If, in addition, the group G is solvable, more information is available about $\chi(1)$. Since $\alpha(1) = 1$, by Exercise 4.12 of [2] we have that $[\chi\psi, \alpha] = 1$. If $\chi\psi(1) \neq 1$, by Proposition 4.3 there exists a character $\beta \in \text{Irr}(G)$ such that $\beta \neq \alpha$ and $[\beta, \chi\psi] = 1$.

The main results of this section are Propositions 4.2 and 4.3. Those are corollaries of Theorem B and C of [1] and Lemma 4.1.

Lemma 4.1. *Let G be a finite group and $\chi\psi \in \text{Irr}(G)$. Set*

$$\chi\psi = \sum_{i=1}^n a_i \alpha_i$$

where $n > 0$ and $a_i > 0$ is the multiplicity of $\alpha_i \in \text{Irr}(G)$ in $\chi\psi$, for each $i = 1, \dots, n$.

If $\alpha_1(1) = 1$, then the irreducible constituents of the character $\chi\overline{\chi}$ are $1_G, \overline{\alpha_1}\alpha_2, \overline{\alpha_1}\alpha_3, \dots, \overline{\alpha_1}\alpha_n$, and

$$\chi\overline{\chi} = a_1 1_G + \sum_{i=2}^n a_i (\overline{\alpha_1}\alpha_i)$$

where $n > 0$ and $a_i > 0$ is the multiplicity of $\overline{\alpha_1}\alpha_i$ in $\chi\overline{\chi}$. In particular, $\eta(\chi\psi) = \eta(\chi\overline{\chi})$.

Proof. Observe that $\alpha_1\overline{\chi} \in \text{Irr}(G)$ since $\alpha_1(1) = 1$ and $\chi \in \text{Irr}(G)$. Since $[\psi, \alpha_1\overline{\chi}] = [\chi\psi, \alpha_1] > 0$, we have that $\psi = \alpha_1\overline{\chi}$.

Since $\psi = \alpha_1\overline{\chi}$ and α_1 is a linear character, we have that $(\alpha_1)^{-1} = \overline{\alpha_1}$ and

$$\chi\overline{\chi} = \sum_{i=1}^n a_i(\overline{\alpha_1}\alpha_i).$$

Observe that $\overline{\alpha_1}\alpha_i \in \text{Irr}(G)$ since α_1 is a linear character and $\alpha_i \in \text{Irr}(G)$. Also observe that $\overline{\alpha_1}\alpha_i \neq \overline{\alpha_1}\alpha_j$ if $\alpha_i \neq \alpha_j$. Thus the distinct irreducible constituents of $\chi\overline{\chi}$ are $1_G = \overline{\alpha_1}\alpha_1, \overline{\alpha_1}\alpha_2, \overline{\alpha_1}\alpha_2, \dots, \overline{\alpha_1}\alpha_n$, and a_i is the multiplicity of $\overline{\alpha_1}\alpha_i$ in $\chi\overline{\chi}$. $\square$

Proposition 4.2. *Let G be a finite solvable group. Let $\chi, \psi \in \text{Irr}(G)$ with $\chi(1) > 1$. If the product $\chi\psi$ has a linear constituent, then $\chi(1) = \psi(1)$ and $\chi(1)$ has at most $\eta(\chi\psi) - 1$ distinct prime divisors.*

If, in addition, G is supersolvable, then $\chi(1)$ is a product of at most $\eta(\chi\psi) - 2$ primes.

Proof. By Theorem C of [1], we have that $\chi(1)$ has at most $\eta(\chi\overline{\chi}) - 1$ distinct prime divisors. Also by Theorem C of [1] if, in addition, G is supersolvable, then $\chi(1)$ is a product of at most $\eta(\chi\overline{\chi}) - 2$ primes. Thus the result follows from Lemma 4.1. $\square$

Proposition 4.3. *Let G be a finite solvable group and $\chi, \psi \in \text{Irr}(G)$ with $\chi(1) > 1$. Assume that the product $\chi\psi$ has a linear constituent α_1 and the decomposition of the character $\chi\psi$ into its distinct irreducible constituents*

$\alpha_1, \alpha_2, \dots, \alpha_n$ has the form

$$\chi^\psi = \sum_{i=1}^n a_i \alpha_i$$

where $n > 0$ and $a_i > 0$ is the multiplicity of α_i for each $i = 1, \dots, n$.

Then $a_1 = [\alpha_1, \chi^\psi] = 1$ and $1 \in \{a_i \mid i = 2, \dots, n\}$.

Proof. Let $\{\theta_i \in \text{Irr}(G)^\# \mid i = 2, \dots, \eta(\chi\overline{\chi})\}$ be the set of non-principal irreducible constituents of $\chi\overline{\chi}$. If $\text{Ker}(\theta_j)$ is maximal under inclusion among the $\text{Ker}(\theta_i)$ for $i = 2, \dots, n$, then $[\chi\overline{\chi}, \theta_j] = 1$ by Theorem C of [1]. Thus $1 \in \{[\chi\overline{\chi}, \theta_i] \mid i = 2, \dots, n\}$. The result then follows from Lemma 4.1. $\square$

References

- [1] E. Adan-Bante, Products of characters and derived length, J. Algebra 266 (2003), 305-319.
- [2] I. M. Isaacs, Character Theory of Finite Groups. New York-San Francisco-London: Academic Press 1976.
- [3] T. M. Keller, Orbit sizes and character degrees III, J. Reine Angew. Math 545 (2002), 1-17.